

Inter-American Drug Abuse Control Commission CICAD




RESULTS OF THE DRUG ABUSE MONITORING SURVEY WITHIN CORRECTIONAL FACILITIES IN BERMUDA

Dr. Kyla Raynor

- Surveillance of drug use and crimes among the inmate population (high risk).
- Classification of levels of severity of substance abuse problems among reception inmates (none, low, intermediate, substantial, severe).
- Needs analysis of mandatory substance abuse treatment identify inmates needing drug treatment while incarcerated (Right Living House).
- Trend analysis of drug consumption and criminality over

DATA COLLECTION (April 1st, 2011 – March 31st, 2012)



- Census of all new inmates in two facilities:
 - Westgate Correctional Facility
 - Co-Ed Facility
- Voluntary

Survey Year	Number of Inmates (Respondents)	Respondents
2004/2005	257	Men only
2006/2007	178	Men only
2011/2012	293	Men, Women, Juveniles

QUESTIONNAIRE

5

Drug Abuse Monitoring Programme:

- DAST-28 (Drug Abuse Screening Test) plus additional items related to drug use history and crime
- Entire questionnaire takes ≈ 30 mins. 2½ hours to complete

Topics include:

- Demographics
- Self Reported Drug Use
- Drug Screening
- Drug Treatment & Perceptions of Need for Treatment
 - n Perceptions of current offences' relationship to drugs and/or alcohol
- Criminal History and Outcomes (current and past offences violent, property, drug, public order, other offences; fine, probation, prison)
- Personal Health & Abuse History (HIV, physically or sexually abused)
- Drug Market
 - n Purchase and distribution of illegal drugs in past year and 30 days
 - n Self-report of other drugs on the streets
- DAST



- Dichotomous response category (Yes/No)
- A score of "1" is given for each YES response, except for items 4, 5, and 7, for which a NO response is given a score of "1"
- Cutoff scores of 6 through 11 are considered to be optimal for screening for substance use disorders
- Using a cutoff score of 6 has been found to provide excellent sensitivity for identifying substance use disorders as well as satisfactory specificity (i.e., identification of persons who do not have substance use disorders).
- using a cutoff score of <11 somewhat reduces the sensitivity for identifying persons with substance use disorders, but more accurately identifies persons who do not have a substance use disorder.
- Over 12 is definitely a substance abuse problem
 - n In a heterogeneous psychiatric patient population, most items have been shown to correlate at least moderately well with the total scale scores.
- The items that correlate poorly with the total scale scores appear to be items 4, 7,16, 20, and 22
- Although DAST-28 was utilised, DADT-20 was used in the analysis and will be used going forward. (none = 0; low = 1 5; intermediate = 6 10; substantial = 11 15; severe = 16 20)


PROFILE OF SURVEY PARTICIPANTS

- Of all inmates within the Department of Corrections, 293 persons responded to the survey.
- 93.2% male offenders; 82.3% were "Black", 72.4% were single; 60.1% with dependents; 46.8% raised by single parents; 39.2% with at most a high school education; 27.6% with part-time or odd jobs; and 54.3% being 35 years or younger.
- Of the 293 persons, 18 (6.1%) were repeat offenders of at most two times each.
- Overall, 99% (n = 290) of the respondents have reported use of at least one drug in their lifetime, and 1 in 5 (19.5%) were poly drug users.

DRUG CONSUMPTION

- Highest lifetime consumption was observed for alcohol (96.9%), marijuana (91.8%), cigarettes/tobacco (87%), crack cocaine (36.5%), and heroin (26.6%).
- Respondents indicated the highest consumption for alcohol (76.8%), cigarettes/tobacco (74.1%), and marijuana (64.2%) in the 30-days prior to the survey.
- Age of initiation of drug use ranges from a low of 14.9 years for alcohol to a high of 31.6 years for methadone.
- Marijuana and cigarette/tobacco use began around 15.1years and the use of heroin and cocaine began after 20 years, on average.

LIFETIME AND CURRENT PREVALENCE



RAPID DRUG SCREEN RESULTS

Rapid Drug Panel	% Positive
THC	50.2
Opiates	11.3
Cocaine	15.4
Methadone	1.7
Ecstasy	1.7
*16.4% of respondents refused drug screening	

Comparison of Self-Reported Current Users of Substances and Positive Urine Tests


(n = 293)

SUBSTANCE		Ported Nt USE	POSITIVE	URINE TEST		USE AND URINE TEST
	n	%	n	%	n	%
Marijuana/THC	188	64.2	147	50.2	132	45.1
Heroin/Opiates	33	11.3	33	11.3	21	7.2
Crack Cocaine	46	15.7	45	1 E /	29	9.9
Cocaine Powder	6	2.0	45	15.4	1	0.3
Ecstasy	5	1.7	5	1.7	3	1.0
Methadone	12	4.1	5	1.7	2	0.7


Overestimation apparent in self-reported use of marijuana/THC; 188 inmates said they had used marijuana in the past 30 days, however urine results suggest fewer, n=147.

Underestimation evident in self-reported use of cocaine powder; 6 inmates indicated use however drug screening results indicate 45 were actually positive for cocaine.


DRUG TREATMENT




CRIMINALITY



DRUG CONNECTION WITH OFFENCE



DRUG MARKET CONNECTION



SELF-REPORTED HISTORY OF ABUSE

TYPE OF ABUSE	%	n
Physical Abuse	16.4	48
Sexual Abuse	5.8	17
Neglect	15.0	44
Abandonment	15.4	44

DAST RESULTS

	Level of Severity	% Respondents
	None (0)	5.5
	Low (1-5)	50.9
	Intermediate (6-10)	21.8
Substance abuse or dependence	Substantial (11-15)	16.4
	Severe (16-20)	5.5

- Overall, 43.7% of reception offenders were classified as having "intermediate to severe" drug abuse problems at the time of their current arrests and incarceration.
- About 84.6% reported that they could not get through the week without using drugs.
- More than half (53.6%) of all offenders indicated their spouse/parents complained about their involvement with drugs.

QUESTIONS?

Thank You

Dr. Kyla Raynor
Senior Research Officer/Policy Analyst
Department for National Drug Control

Government of Bermuda



"Attempts to deter drug use through punishment fail because they do not address the complex causes of drug abuse, which begins within the context of family problems and peer deviant behavior."

United States National Institute of