

### Contaminated Sites in Nicaragua

PLACE	PTS	DESTINY	AMOUNT
León	DDT	water	
León	Aldrin, Endrin	water	
León (Aerop. Godoy)	Toxaphene	water	25-35 mg/l
León	Parathion	water	
Managua	DDT	water	
Managua	Aldrin, Endrin	water	
Managua (Lago)	Toxaphene	water	0.001-0.008 ppm
Managua	Parathion	water	
lago Xolotlán	Toxaphene	animals	0.15 - 19.5 mg/kg
Chinandega	Toxaphene	water	0.2-25 ppb / 12 mg/l
Chinandega	Toxaphene	animals	
Chinandega	Toxaphene	humans	
Chinandega	Aldrin, Endrin	humans	
Chinandega	DDT	humans	
Chinandega	DDT	water	
Chinandega	Parathion	water	
Chinandega	Aldrin, Endrin	water	
El Picacho	Toxaphene	water	18 mg/l
El Picacho	Toxaphene	soil	5600 mg/kg
El Picacho	DDT	soil	
Chinandega	Toxaphene	soil	16 mg/kg
Sébaco	Toxaphene	soil	9 mg/kg
río Atoya	Heptachlor	water	7.5 mg/kg
río Sasama	Heptachlor	water	7.5 mg/kg
estero Naranjo	Toxaphene	animals	
Estero Naranjo	DDT	animals	
río Atoya	Heptachlor	humans	
León (Aerop. Godoy)	Toxaphene	soil	3500 mg/kg
León (Aerop. Godoy)	DDT	soil	
León (Aerop. Godoy)	Endosulfan	soil	
El Picacho	various	air	
Paz Centro, León	Toxaphene	water	12 mg/l
Telica, León	Toxaphene	water	0.1-0.6 mg/l
Managua (Lago)	Toxaphene	sediment	0.09-1.4 ppm

\*SOURCE: Sitios Contaminados con COP en Nicaragua, MARENA, 2004  
© 2007 OAS